

2022 SCOTT RIVER SALMON STUDIES FINAL REPORT



Prepared By:

Morgan Knechtle and Domenic Giudice

California Department of Fish and Wildlife
Northern Region
Klamath River Project
1625 South Main Street
Yreka, CA 96097

2022 SCOTT RIVER SALMON STUDIES

California Department of Fish and Wildlife
Northern Region
Klamath River Project

ABSTRACT

The California Department of Fish and Wildlife's (Department) Klamath River Project (KRP) operated an adult fish counting facility and conducted cooperative spawning ground surveys (carcass surveys) on the Scott River during the 2022 fall-run Chinook Salmon (*Oncorhynchus tshawytscha*) and Coho Salmon (*Oncorhynchus kisutch*) spawning season. The purpose of these surveys is to describe the run characteristics of adult Chinook Salmon and Coho Salmon into the Scott River. Fish counting operations began on September 29, 2022 and ended on December 26, 2022. Incidental monitoring of steelhead (*Oncorhynchus mykiss*) also occurs.

The total number of Chinook Salmon that entered the Scott River during the 2022 season is estimated to be **994** fish. Based on the proportion of male and female Chinook Salmon that were sampled during the spawning ground surveys, the run was comprised of approximately 309 (31.1%) males and 685 (68.9%) females. Based on scale age analysis, adults comprised approximately 93% (927 fish) and grilse comprised 7% (67 fish) of the run. Males ranged in fork length (FL) from 41 cm to 91 cm and averaged 68.1 cm. Females ranged in FL from 42 cm to 85 cm and averaged 66.5 cm. KRP staff estimated that none of the Chinook Salmon that returned were of hatchery origin.

The first adult Coho Salmon was observed at the Scott River Fish Counting Facility (SRFCF) on November 30, 2022, and the last Coho Salmon was observed on December 26, 2022. A net total of **238** Coho Salmon were estimated in the Scott River during the season. The counting station was removed during the day with the highest observed daily Coho Salmon migration. If the counting station was not removed additional Coho Salmon would have been counted. Based on live Coho Salmon images observed during video monitoring, adults comprised approximately 97.1% and grilse comprised 2.9% of the run. Five Coho Salmon carcasses were recovered opportunistically by the Siskiyou Resource Conservation District. Based on observed hatchery marks during the carcass survey and inspections of images during the video survey, none of the Coho Salmon were estimated to be of hatchery origin.

The first steelhead >40.6 cm was observed on November 6, 2022, and the last steelhead was observed on December 26, 2022. During this time there were a total of 26 steelhead observations including both upstream (22) and downstream (4) movements. The net upstream movement of **18** fish represents a minimum number of steelhead for the season. A large fraction of the adult steelhead migration occurs outside the operational window of the Scott River Fish Counting Facility. Zero steelhead were recovered during the spawning ground survey effort.

INTRODUCTION

STUDY LOCATION AND RUN TIMING

The Scott River is a major tributary of the Klamath River located in Siskiyou County, and enters the Klamath River at river mile 143 (Figure 1). The Scott River Fish Counting Facility (SRFCF) is located at river mile 18.2, (041° 38' 10.93" N; 123° 04' 3.08"W) 0.63 river miles downstream of the Jones Beach picnic area near the transition between the canyon and valley reaches. Chinook Salmon typically return to the Scott River to spawn from mid-September through late December. Coho Salmon typically return to spawn from mid-October through early January and steelhead from November through April.




Figure 1. Location of the Scott River, tributary to the Klamath River, Siskiyou County, California.

KLAMATH RIVER PROJECT AND THE SCOTT RIVER STUDY

The Scott River study is one component of the Klamath River Project (KRP) (initiated in 1978). The goals of the KRP include obtaining information on species composition, hatchery composition, run timing, age structure, spawning distribution, fork length (FL) frequency and sex ratios in various tributaries to the Klamath River including the Salmon, Scott, and Shasta rivers, as well as Bogus Creek and 22 other smaller tributaries. The Scott River is particularly important because it is a major salmon spawning tributary. In addition to providing valuable escapement estimates to the Pacific Fisheries Management Council for the effective management and allocation of fall Chinook Salmon originating from the Klamath River Basin, the Scott River studies provide an opportunity to monitor an independent population of Coho Salmon (Williams et al. 2008) within the state and federally threatened Southern Oregon/Northern California Coast Coho (SONCC) range. The Scott River Coho Salmon population is currently believed to be the largest sub population of wild Coho Salmon within the Klamath watershed.

In the early years of the KRP, spawning ground surveys were conducted in the major spawning areas of the main stem Scott River which included an approximately 5.5 mile reach near Etna and a 4.75 mile reach downstream of the State Highway 3 Bridge crossing near Fort Jones. From 1989 through 1991 spawning ground surveys were limited to the lower river. In 1985 a temporary fish marking weir was installed on the lower river at river mile 1.6 and was operated during each spawning season until 1991 (Figure 2). Operation of the weir was often hampered by high flows and beginning in 1992 operation of the marking weir was dropped in favor of conducting more intensive mark recapture spawning ground surveys in cooperation with United States Forest Service (USFS) fisheries staff. Currently the spawning ground survey is a cooperative effort between the USFS, Quartz Valley Tribe, Karuk Tribe, Northern California Resource Center, Siskiyou Resource Conservation District, Siskiyou County Schools and the Department.




Figure 2. Chinook Salmon observed in the lower Scott River (river mile 1.6) during trapping efforts from 1985-1991.

In 1994 the California State Legislature passed the Leslie Amendment (SB 779). The passage of SB 779 required Departmental staff to obtain landowner permission prior to accessing private lands to conduct biological investigations. As a result, since 1994, spawning ground surveys have been limited to those areas of the river on private land where landowner permission has been granted. The entire length of the Scott River within the Scott Valley passes through private ownership. The level of cooperation from local landowners has varied over the years. Controversies associated with the listing of SONCC Coho Salmon under the California Endangered Species Act (CESA) and other regulatory actions have reduced the amount of cooperation provided by local landowners to the extent that the Department has been denied permission to survey a large portion of the salmon spawning reaches present in Scott Valley. As a result of the limited landowner access to the valley reaches, since 2007 the Department has been operating the Scott River Fish Counting Facility (SRFCF) at the upper end of the canyon. The addition of the SRFCF has allowed for an accurate estimation of salmonid migration to areas of spawning habitat in the valley reaches without having to conduct spawning ground surveys in these reaches. Since 2014 the Department has exercised its authority under the navigability statute to access the Scott River adjacent to private lands in the lower 18.2 river miles (downstream of the counting facility). The location of the fish counting station allows for monitoring fish abundance into the valley while Cormack-Jolly-Seber mark recapture carcass-based or redd based estimates are conducted in the areas downstream of the counting station. The counting facility is also located upstream of several tributaries that can produce significant fall and winter stream flows, thereby reducing the probability of having the counting facility inoperable due to high flow events.

SCOTT RIVER STUDY OBJECTIVES SUMMARIZED:

- Determine the in-river run size (escapement) of Chinook Salmon and Coho Salmon returning to the Scott River.
- Determine run timing, spawning distribution, length frequency (FL) distribution, and sex ratio for Chinook Salmon and Coho Salmon in the Scott River.
- Collect scale samples from carcasses and look for hatchery marks to determine age composition and hatchery contribution rates of the run.
- Collect biological data for all steelhead observed coincidentally during the Chinook Salmon and Coho Salmon spawning seasons.

METHODS

OPERATION OF THE SCOTT RIVER FISH COUNTING FACILITY

The video fish counting system was installed at the SRFCF on September 29, 2022, at 1200 hours Pacific Standard Time (PST). A temporary weir (Alaskan weir design) was installed to direct migrating fish into a flume where they pass in front of a video camera (Figure 3). The underwater video system consisted of a digital video camera, waterproof camera housing, viewing window, and counting flume which allowed for recording unimpeded fish passage through the facility. The facility was operated 24 hours a day, seven days a week during the

Chinook Salmon and Coho Salmon migration. A Splash Cam digital video camera equipped with a 3.6mm wide angle lens with an auto iris was used to collect the video image and an Ever Focus Digital Video Recorder (Model ECOR 264) was used to record the image to external hard drives. The time lapse DVR was set to record continuously, and data storage drive changes were made at least twice a week.



Figure 3. Scott River Fish Counting Facility located in Siskiyou County, California 2022.

All hard drives were immediately returned to the office where each was subsequently downloaded and reviewed by project staff in the video lab. During each review staff recorded the date, time (hour:min:sec), and species of each fish observed on each video image. Additionally, salmon are preliminarily classified as either adults or grilse based on observed length in the videography. During videography adult salmon are considered greater than and equal to 56 cm and salmon less than 56 cm are considered grilse. All rainbow trout (*onchorhynchus mykiss*) observed during videography greater than 40.6 cm (16 inches) in length are considered adult steelhead. If the species could not be determined due to poor visibility or picture quality, staff recorded that observation as species unknown. Staff also noted any adipose fin-clipped or maxillary-clipped fish observed and recorded the presence of lamprey and any other distinguishable marks that were visible on the image. All data was then entered into computer files and each data file was subjected to one independent review prior to commencement of data analysis.

SPAWNING GROUND SURVEYS

Spawning ground surveys on the Scott River main stem were conducted twice a week on Mondays and Thursdays and opportunistically in main stem tributaries downstream of the counting facilities throughout the salmon spawning season starting October 10, 2022 and

ending December 12, 2022. Additional surveys were conducted upstream of the counting facility in the main stem and select tributaries during the Coho Salmon period and are reported by the Siskiyou Resource Conservation District (SRCD). A total of 80 surveys were performed during the spawning season (Appendix 1). On the morning of each survey, crews of at least two people each were given daily instructions, data sheets, field equipment, vehicle assignments, and were assigned a survey reach. Crews walked their assigned reach in a downstream direction looking for salmon carcasses and spawning redds. All new redd locations were flagged, mapped on USGS topographic maps, and geo referenced using GPS coordinates. This information was provided to the Klamath National Forest for use in their annual reporting. All carcasses recovered were identified to species and gender, checked for marks or tags, and measured (FL); a scale sample was collected for age composition analysis, and females were examined for spawning success.

For purposes of the mark recapture estimate, each carcass was categorized into one of four pathways (Path). Fresh carcasses, those with clear eyes and/or firm flesh were designated as Path 1. Individually numbered jaw tags were attached to the lower right jaw of all Path 1 carcasses and returned to the river for potential recapture during later surveys. Older carcasses, those with two cloudy eyes and/or mushy flesh, were categorized as Path 2. All Path 2 carcasses were cut in half and returned to the river after all biological data was collected. Path 3 carcasses included all of the Path 1 carcasses (with jaw tag) that were recaptured during subsequent surveys. Path 3 carcasses were returned to the river for future recapture if the adipose fin clip determination could still be made with confidence. Therefore Path 3 carcasses could be recaptured multiple times. Once an adipose fin had deteriorated to the point that adipose fin clip determination could not be made with confidence, the survey tag was removed and the carcass was chopped in half and removed from the mark recapture experiment. Any carcasses that could be observed by a survey crew but could not be retrieved for data collection, because they were located in inaccessible or unsafe locations, were designated as Path 4.

During 2022 the Cormack-Jolly-Seber (CJS) model (Bergman et al. 2012) was utilized to estimate Chinook salmon abundance in the area downstream of the counting station.

SURVEY REACHES

Survey reaches have remained consistent since the beginning of the cooperative spawning ground survey in 1992. During the Chinook Salmon spawning season, decisions regarding which reaches should be surveyed were based on the known distribution of the Chinook Salmon run each week, the available labor force present during each survey, and reach specific stream conditions.

A total of 16 survey reaches, covering approximately 53.6 river miles, have been identified on the Scott River (Table 1, Figure 4). Historically, the highest observed densities of Chinook Salmon spawning areas within Scott Valley were located downstream of the State Highway 3 Bridge crossing (river mile 34.6) to the USGS gauging station located at river mile 21 (Reaches 8, 9, and 10), and in that part of the river located downstream of Young's Dam, (river mile 46 to about river mile 42) located upstream of the Eller Lane Bridge crossing (Reaches 12-14).

Table 1. Description of cooperative spawning ground survey reach locations along the Scott River during the 2022 season.

Reach Number	Downstream Limit	RM	Upstream limit	RM	Length (miles)
1	Mouth	0.00	Mid Point	2.55	2.55
2	Mid Point	2.55	Pat Ford	5.14	2.59
3	Pat Ford	5.14	George Allen Gulch	8.51	3.37
4	George Allen Gulch	8.51	Gold Flat	11.38	2.87
5	Gold Flat	11.38	Bridge Flat	14.73	3.35
6	Bridge Flat	14.73	Counting Weir	18.20	3.47
7	Counting Weir	18.20	USGS Stream Gage	21.66	3.46
8	USGS Stream Gage	21.66	Meamber Bridge	25.36	3.70
9	Meamber Bridge	25.36	Dunlop	29.50	4.14
10	Dunlop	29.50	Highway 3 Bridge	35.60	6.10
11	Highway 3 Bridge	35.60	Eller Lane	41.10	5.50
12	Eller Lane	41.10	Sweezy Bridge	42.10	1.00
13	Sweezy Bridge	42.10	Horn Lane	43.90	1.80
14	Horn Lane	43.90	Young's Dam	46.00	2.10
15	Young's Dam	46.00	Fay Lane	49.60	3.60
16	Fay Lane	49.60	East Fork Confluence	53.60	4.00




Figure 4. Location of the Scott River Fish Counting Facility and mainstem spawning ground survey reaches on the Scott River used during the 2022 field season.

To assist in developing stock identification baseline information, the KRP collected both genetic tissue and otolith samples during the season. Tissue samples were collected for future DNA analysis from 43 Chinook Salmon and five (RCD collection) Coho Salmon. Tissue was collected from the first Chinook Salmon from each reach and each survey date and all Coho Salmon for which samples could be collected. All samples were collected following protocols provided by the National Oceanic Atmospheric Administration’s (NOAA) Southwest Fisheries

Science Center. Samples were sent to the Salmonid Genetic Tissue Repository located at the NOAA Santa Cruz Laboratory for archiving and analysis. Otoliths were collected from 39 Chinook Salmon and five (RCD collection) Coho Salmon. Otoliths were collected from the first Chinook Salmon from each reach and each survey date and all Coho Salmon for which samples could be collected. All otoliths collected were archived for future microchemistry analysis. All otolith samples were collected following standard protocols described by Stevenson (1992).

POPULATION ESTIMATES

The Chinook Salmon spawner escapement for the Scott River upstream of the counting facility was derived from a direct count of all Chinook Salmon observed at the counting facility (upstream minus downstream observations). To estimate adult escapement in the Scott River downstream of the counting station, the number of Chinook salmon carcasses derived from the Cormack-Jolly-Seber model were estimated (utilizing data from Reach 1 through Reach 6 only). To estimate total Chinook escapement, the number of Chinook estimated downstream of the counting facility was summed with the estimate from upstream of the counting station.

The Coho Salmon spawner escapement for the area of the Scott River upstream of the counting facility was also derived from a direct count of all Coho Salmon observed at the counting facility. Spawning ground surveys were conducted through December 14, 2022 in the main stem and tributaries (Tompkins Creek, Kelsey Creek, and Canyon Creek) below the counting facility. To estimate total adult Coho Salmon escapement in the Scott River, the number of observed Coho Salmon redds downstream of the counting station (zero in 2022) were multiplied by two in order to estimate the number of adult Coho Salmon (assuming two unique individuals participated in the construction of each redd) and were added to the count of all Coho Salmon observed passing through the SRFCF. The grilse component below the counting facility was estimated using the following equation: $\text{total run} = \text{adults} / (1 - \% \text{grilse})$. Zero Coho Salmon redds were identified downstream of the counting station during 2022. Therefore, no Coho Salmon were estimated downstream of the counting station.

HATCHERY CONTRIBUTION RATES

The hatchery contribution rates for Chinook Salmon and Coho Salmon have been estimated both through the recovery of carcasses and through reviewing live fish images observed at the fish counting facility. During the 2022 season hatchery contribution rates were based on observed clips on recovered carcasses for both Chinook Salmon and Coho Salmon. The hatchery contribution rate of Chinook Salmon was calculated by multiplying the number of CWTs observed for each CWT group by its production multiplier value (the inverse of the proportion of each group of juveniles that were tagged).

An additional sample expansion (the inverse of the number of fish handled during spawning ground surveys divided by the total estimated) was applied. The Coho Salmon hatchery contribution rate was estimated by direct observation from Coho Salmon carcasses recovered during the spawning ground survey. The observed clip rate was then applied to the total estimated Coho Salmon run size.

RESULTS

OPERATION OF THE SCOTT RIVER FISH COUNTING FACILITY

The SRFCF began recording fish movement on September 29, 2022. The first Chinook Salmon was observed at the SRFCF on October 15, 2022, and the last Chinook Salmon was observed on December 7, 2022. The run peaked between October 26, 2022 and November 6, 2022 when 82.4% of the Chinook Salmon migration was observed (Figure 5). Unlike in previous years when over 90% of Chinook Salmon passed through the SRFCF during daylight hours, in 2019, 2020, 2021 and 2022 55%, 64%, 63% and 39% of the Chinook Salmon were observed during daylight hours (Figure 6).




Figure 5. Run timing of Chinook Salmon through the Scott River Fish Counting Facility during the 2022 season (N=74), and average daily flows observed at USGS Gauge No. 11519500.

A net total of 74 Chinook Salmon were estimated to have passed through the SRFCF during the 2022 season (76 upstream and 6 downstream) when videography was employed. During the season the video monitoring station operated continuously, and no video malfunctions occurred. After accounting for upstream known and unknown observations a seasonal estimate of **74** Chinook Salmon upstream of the counting station was generated.




Figure 6. Summary of daily run timing of Chinook Salmon observed at the Scott River Fish Counting Facility during 2022 (N=74).

SPAWNING GROUND SURVEYS

A total of 212 Chinook Salmon were encountered in all areas during the spawning ground survey as Path 1 or Path 2 carcasses. Of these, 206 were identified as Path 1 or Path 2 carcasses of which 64 (31.1%) were male and 142 (68.9%) were female (two unsexed carcasses were measured and 4 carcasses were sampled but not measured). Males ranged in FL from 41 cm to 91 cm and averaged 68.1 cm (Figure 7). Females ranged in FL from 42 cm to 85 cm and averaged 66.5 cm (Figure 8). No ad-clipped Chinook were observed during the spawning ground survey effort. After examination of the length frequency distribution of Path 1 and Path 2 carcasses, a preliminary maximum grilse cut-off of 57 cm was established for Scott River Chinook Salmon. The preliminary grilse cut-off was supported by the final age determination which was derived from aged scales.




Figure 7. Length Frequency distribution of Path 1 and Path 2 male Chinook Salmon observed during spawning ground surveys in the Scott River, 2022 (N=64).




Figure 8. Length frequency distribution of Path 1 and Path 2 female Chinook Salmon observed during spawning ground surveys in the Scott River, 2022 (N=142)

Each female carcass was examined to determine if it had successfully spawned prior to death. Spawning status was defined as un-spawned (many eggs remaining in the body) or spawned (few or no eggs remaining). Of the 142 female Chinook Salmon carcasses examined for spawning status, 136 females (95.8%) were found to have spawned, and 6 female (4.2%) carcasses were identified as un-spawned.

The Cormack-Jolly-Seber (CJS) mark recapture method was used to estimate carcass abundance in the area downstream of the counting station. A total of 920 (+/- 95%CI) Chinook Salmon carcasses were estimated (Table 2) below the counting facility utilizing the CJS method (212 carcasses encountered; 172 carcasses marked and released; 34 recapture events from 29 carcasses).

Table 2. Chinook Salmon grilse, adult and total estimates for upstream and downstream of the counting station and total estimates for the Scott River 2022.

	Estimated Chinook Upstream of Counting Station	Estimated Chinook Downstream of Counting Station		Total Chinook Estimates for Entire River	
	Video	CJS	Redds	CJS + Video	Redds + Video
Grilse	5	64	54	70	59
Adults	69	856	714	924	783
Total	74	920	768	994	842

Alternatively, Chinook Salmon abundance downstream of the counting station was estimated through expanding the seasonal total redd count of 357 redds. Multiplying the seasonal total redd count by 2 generates an adult estimate of 714. 54 grilse were added to the adult estimate utilizing the following equation: total run= adults/(1-%jacks estimated). A scale-based grilse estimate of 7.0% was used from recovered carcasses during the spawning ground survey. Utilizing the redd estimate with the grilse proportion yields an estimated total of **768** Chinook Salmon downstream of the counting station.

There is a discrepancy between the CJS (920) and redd based (768) estimate downstream of the counting station of 152 Chinook Salmon (Table 2). We elected to use the carcass based CJS model to estimate the Chinook population downstream of the counting station. The CJS estimate is our preferred method to estimate this sector and we have deferred to the redd estimate only when there is gross disagreement between the two estimates.

To estimate the total Chinook Salmon run for the entire river the CJS estimate from downstream of the counting station (920) was added to the video estimate (74) from upstream of the counting station to yield a total basin run size estimate of **994**. Based on scale age analysis, adults comprised approximately 93.0% (924) and grilse comprised 7.0% (70) of the run (KRTAT 2023).

COHO SALMON

The first adult Coho Salmon was observed at the counting facility on November 30, 2022 and the last Coho Salmon was observed on December 26, 2022 (Figure 9). The counting station was removed for the season on December 26, 2022 due to high flows. A net total of 238 Coho Salmon (239 upstream and 1 downstream) were observed moving through the SRFCE.

During the season the video monitoring station operated continuously, and no video malfunctions occurred.




Figure 9. Run timing of Coho Salmon observed passing through the Scott River Fish Counting Facility during the 2022 season (N=238), and average daily flows observed at USGS Gauge No. 11519500.

Coho Salmon migration peaked on two occasions during the 2022 season. Both of the pulses of migration were associated with increases in base flow. These periods of migration were observed from December 13, 2022 through December 15, 2022, and from December 23, 2022 through December 26, 2022. During these seven days 97.5% of the Coho Salmon (232) were observed. The counting station was removed during the day with the highest observed daily Coho Salmon migration. If the counting station was not removed additional Coho Salmon would have been counted. During the 2022 season KRP staff estimated the number of grilse in the Scott River by enumerating the number of Coho Salmon observed in the video flume that were shorter or longer than 56 cm. Utilizing this method KRP staff identified 97.1% adults and 2.9% grilse (see below for further information on seasonal estimates of age proportions).

Diel movements of Coho Salmon through the SRFCF were consistent throughout the darkest hours of the day and early morning, with fewer observations from 10:00 am and 4:00 pm (Figure 10). In recent years of Coho Salmon monitoring, migrations were generally low during the day and increased from the late afternoon through early morning.




Figure 10. Diel migration patterns of Coho Salmon observed moving through the Scott River Fish Counting Facility in 2022 (N=238).

SPAWNING GROUND SURVEYS

No Coho Salmon carcasses were observed during the cooperative spawning ground surveys on the lower mainstem Scott River (Reaches 1-8) and 5 Coho Salmon carcass were observed opportunistically in upper Scott River (Reaches 9 through 16 and tributary reaches) by the Siskiyou Resource Conservation District (Voight, 2023 pers comm). All the Coho Salmon carcasses recovered were males ranging in fork length from 58 cm to 73 cm (Figure 11). None of the recovered Coho Salmon carcasses were determined to have had clips or marks associated with hatchery production resulting in an estimated hatchery proportion of zero.

Applying the preliminary maximum grilse fork length of 56 cm to the total number of measured carcass recoveries (5) generated an age two proportion of 0% and an age three proportion of 100% (see below for further information on seasonal estimates of age proportions). All 5 of the carcasses examined were sampled for scales and tissue. Collected tissue samples were supplied to the NOAA Southwest Fisheries Science Center located in Santa Cruz, California for stock identification purposes. Additionally, 5 Coho Salmon otolith samples were collected and archived at the Department’s Yreka office for future stock identification and life history examination. Zero Coho Salmon redds was observed during surveys in Kelsey Creek, Canyon Creek or Tompkins Creek.




Figure 11. Length frequency distribution of Male Coho Salmon observed (N=5) during the cooperative spawning ground survey (0) and the SRCD (Voight 2023) efforts (5) in the Scott River 2022-2023.

A total of 238 Coho Salmon were estimated to have swum upstream through the SRFCF during the monitoring period. The age two proportions generated from videography and recovered carcasses were 2.9% and 0%, respectively. Due to the larger sample size available to estimate the age two proportion in the videography (238) versus measured carcass recoveries (5), the final age two proportion was determined using the videography data. Therefore, the resulting number of age two and three Coho Salmon are 7 (2.9%) and 231 (97.1%), respectively.

STEELHEAD

In 2022, a net total of 18 adult (>40.64 cm) steelhead (Figure 12) were estimated to have entered and remained in the Scott River during the video recording season from September 29, 2022 to December 26, 2022. Lines on the back of the video flume were set at 40.64 cm (16 inches) to delineate sub-adults versus adults. The 2022 season was the thirteenth monitoring season that lines delineating adult steelhead and sub-adult steelhead were used.




Figure 12. Run timing of steelhead trout (>40.64 cm) observed passing through the Scott River Fish Counting Facility during the 2022 season (N=18), and average daily flows observed at USGS Gauge No. 11519500.

DISCUSSION

CHINOOK SALMON RUNS

Since 1978, the Chinook Salmon run in the Scott River has ranged from 14,477 fish (1995) to 467 fish (2004) and has averaged 4,821 fish (Figure 13). The 2022 Chinook Salmon run in the Scott River ranks 42 (994 fish) out of 45 years of monitoring. The 2022 run was 79% lower than the 45-year average of 4,821 Chinook Salmon. Chinook Salmon escapement to the Scott River from 2015 to 2022 averaged 1,673 fish, a 65% reduction from the historical average (4,921). Average escapement for the Klamath Basin from 2015-2021 is also down from the historical average, by 43% (CDFW 2022). It is concerning that the Scott River Chinook Salmon population is decreasing at a faster rate than the Klamath Basin as a whole.




Figure 13. Estimated escapement of Chinook Salmon returning to the Scott River from 1978 to 2022.

The various proportions of the Chinook Salmon distribution for years 2008-2022 are detailed in Table 3 and identifies the importance of the entire Scott River watershed to Chinook Salmon. From 2008 through 2022 an average of 62% of the Chinook Salmon run spawned upstream of the counting station. Adult Chinook Salmon migration was impeded by low flows in 2022 and an estimated 7% of the Chinook Salmon run was observed passing through the counting station. It is important to track this metric as it helps describe the spatial distribution of annual spawning. There is a lower risk of catastrophic loss due to potential redd scour when eggs are deposited throughout the watershed. Due to low fall flows more than 50% of the Chinook Salmon spawning occurred in canyon reaches in the 2015, 2018, 2020 and 2022 spawning seasons.

Table 3. Scott River Chinook Salmon abundance estimates by area and percentages of the total above and below the Counting Station during the 2008-2022 seasons.

Year	Downstream of Counting Station	Upstream of Counting Station	% Downstream of Counting Station	% Upstream of Counting Station	Total Basin Estimate
2008	1,439	3,234	31%	69%	4,673
2009	1,014	1,197	46%	54%	2,211
2010	280	2,228	11%	89%	2,508
2011	983	4,538	18%	82%	5,521
2012	1,208	8,144	13%	87%	9,352
2013	1,252	3,372	27%	73%	4,624
2014	2,995	9,476	24%	76%	12,471
2015	1,741	372	82%	18%	2,113
2016	363	1,152	24%	76%	1,515
2017	297	2,279	12%	88%	2,576
2018	875	404	68%	32%	1,279

2019	537	1,553	26%	74%	2,090
2020	586	269	69%	31%	855
2021	561	1,400	29%	71%	1,961
2022	920	74	93%	7%	994
Average	1,003	2,646	38%	62%	3,650

The timing of Chinook Salmon passage through the SRFCF has consistently started in early October. Except for 2020, the Chinook Salmon run has migrated through the SRFCF almost entirely during October. During 2022, 82% of the cumulative annual migration occurred in a narrow 12-day period between October 26 and November 6 (Figure 14). The years 2015, 2018, 2020 and 2022 were the four driest falls during the period of monitoring at the counting facility. The proportion of Chinook Salmon that spawned downstream of the counting station in 2015, 2018, 2020 and 2022 were 82%, 68%, 69% and 93% respectively which corresponded with the four lowest average October flow years (CDFW 2021).




Figure 14. Cumulative percent of total Chinook Salmon observations by day at the Scott River Fish Counting Facility annually from 2008-2022. The date in parenthesis indicates the last date the fish counting facility was operated for each year.

The Scott River is an important component of the Klamath Basin Chinook Salmon runs. The Scott River has contributed an average of 9% of the basin-wide (including Trinity River) natural spawning escapement to the Klamath River basin during the period from 1978 to 2022 (Table 4). In most years the Scott River Chinook Salmon population tracks very similarly to the total Klamath Basin population ($r=0.7793$ $p\text{-value} < 0.01$) indicating that environmental factors that affect the entire Klamath basin play an important role in influencing abundance of this population (Figure 15). While this relationship is significant over the period of record the relationship in the most recent 8 years is not significant ($r=-.0516$ $p\text{-value} < .1$).

Table 4. Klamath Basin and Scott River Chinook Salmon natural spawner escapements (age 2-5), 1978-2022.

Year	Chinook Natural Spawner Escapement	% Scott
------	------------------------------------	---------

	Klamath Basin	Scott River	
1978	74,906	5,332	7%
1979	37,398	3,824	10%
1980	48,465	4,277	9%
1981	50,364	6,556	13%
1982	50,597	10,176	20%
1983	33,310	3,568	11%
1984	21,349	1,801	8%
1985	61,628	4,408	7%
1986	142,302	8,041	6%
1987	110,489	8,566	8%
1988	91,930	5,200	6%
1989	49,377	4,188	8%
1990	16,946	1,615	10%
1991	12,367	2,165	18%
1992	17,171	2,838	17%
1993	25,683	5,300	21%
1994	38,578	2,863	7%
1995	179,118	14,477	8%
1996	87,500	12,097	14%
1997	50,369	8,561	17%
1998	45,343	3,327	7%
1999	28,904	3,584	12%
2000	89,122	6,253	7%
2001	85,581	6,142	7%
2002	69,502	4,308	6%
2003	89,744	12,053	13%
2004	28,516	467	2%
2005	27,931	756	3%
2006	45,002	4,960	11%
2007	61,741	4,505	7%
2008	48,073	4,673	10%
2009	52,702	2,211	4%
2010	49,027	2,508	5%
2011	110,554	5,521	5%
2012	137,724	9,352	7%
2013	69,986	4,624	7%
2014	112,599	12,470	11%
2015	31,607	2,113	7%
2016	15,818	1,515	10%
2017	35,036	2,576	7%
2018	61,561	1,279	2%
2019	26,412	2,090	8%
2020	31,138	855	3%
2021	36,564	1,961	5%
2022	26,201	994	4%
Average	58,139	4,821	9%




Figure 15. Chinook Salmon Klamath Basin natural spawner escapement (primary y-axis) and the Scott River natural spawner escapement (secondary y-axis) from 1978 through 2022.

To evaluate freshwater productivity, the production of emigrating 0+ Chinook Salmon has been estimated in the Scott River since Brood Year 1999 (Morrow et al. 2022). The number of 0+ Chinook Salmon produced per adult has been calculated for Brood Years 1999-2015, 2017-2021 and has ranged from a low of 17.6 to a high of 453.6 and averaged 137 (Figure 16). Estimates of 0+ Chinook Salmon recruits per adult for Brood Year 2021 was 251.4. The observed 0+ Chinook Salmon produced per adult for Brood Year 2021 was 83% higher than the 22-year average of 137.7 fish. Similarly, when the number of returning adults is plotted against the number of 0+ Chinook produced a positive relationship is generated (Figure 17). This positive relationship indicates that as adult spawners increase so does the resulting offspring. It is theorized that the extremely low production of offspring from the 2005 brood year was the result of egg scour when a significantly high proportion of the Chinook redds were constructed in the canyon reaches and stream flows during incubation exceeded 23,000 cfs on December 31, 2005. It is theorized that a large fraction of the 2005 Chinook cohort was destroyed during these storms.




Figure 16. Number of 0+ Chinook Salmon produced per adult spawner in the Scott River by brood year, for Brood Years 1999-2015, 2017-2021.

Due to funding constraints the Department was unable to operate the Scott River Rotary Screw Trap during the 2017 season preventing this analysis for Brood Year 2016. As the watershed approaches carrying capacity the number of 0+ Chinook Salmon produced per adult is a direct measure of in-river productivity, and as habitat conditions improve or diminish, this measure will reflect those conditions.




Figure 17. Number of returning adults and corresponding number of 0+ Chinook Salmon produced by year in the Scott River by brood year, for Brood Years 1999-2015, 2017-2021.

COHO SALMON

Since video operations began in 2007 the estimated escapement of Coho Salmon in the Scott River has ranged from a low of 63 to a high of 2,752 and averaged 703 (Figure 18). The adult run size of Coho Salmon prior to 2007 is unknown, and with the addition of the counting facility, the ability to monitor this ESA listed run has greatly improved. Although recent adult run size data is sparse on the Scott River, monitoring of the yearling juvenile emigration has taken place since 2003. The emigration data generated from 2003 through 2022 indicates significant interannual variation (Morrow et al. 2022). Results of the adult monitoring at the SRFCF support this observation. The brood year that returned in 2022 (age 3 in 2022 and age 2 in 2021) in one generation decreased in total numbers by 127 fish to 238, a decrease of 34.8 percent, from returns in 2019 (Figure 19). In 5 generations this brood year has decreased from an estimated 1,622 in 2007 to 238 in 2022.




Figure 18. Estimated escapement by return year of adult and grilse Coho Salmon (age 2 and age 3) returning to the Scott River from 2007 to 2022.

The 2022 monitoring season extended to December 26th near what is considered the normal end of the Coho Salmon migration. The counting facility was removed during active Coho Salmon movement, and it is unknown how many fish were missed after the counting station was removed. To evaluate the accuracy of the 2022 estimate, the proportion of the total annual run that has been observed by December 26th each season for the period of record (Figure 20) has been averaged. For the years 2007-2021, an average of 96.8% of the annual Coho Salmon run has been observed by December 26th. If 96.8% of the Coho Salmon were observed during the 2022 season, then an estimated 8 additional Coho Salmon (total 246) would have been estimated if monitoring was conducted through the entire migration based on previous run timing observations. Although this analysis provides evidence that if adults were missed at the end of the migratory window due to removing the counting facility it was likely a small fraction, it is unknown how many were missed. If the counting station had not removed additional Coho Salmon would likely have been counted.




Figure 19. Estimated escapement by Brood Year of adult and grilse Coho Salmon (age 2 [year-1] and age 3 [year]) salmon returning to the Scott River from 2007 to 2022. Individual brood years are represented by different colors.




Figure 20. Cumulative percent of total Coho Salmon observations by day at the Scott River Fish Counting Facility annually from 2007-2022. The date in parenthesis indicates the last date the fish counting facility was operated for each year.

Utilizing the number of Coho Salmon smolts produced in the Scott River (Morrow et al. 2022) and the results of the adult abundance estimates allows for analysis of Scott River freshwater production and out of basin survival by brood year. For Brood Years 2004-2008, 2010-2014, and 2016-2019, the out of basin survival has averaged 6.1% and ranged from a low of 1.25% to a high of 13.51% (Table 5). Due to the extremely high observed percent smolt survival of 55.81 for Brood Year 2009, data from this brood year has been omitted from this analysis. It is possible that the smolt estimate generated for Brood Year 2009 underestimated the actual number of out migrants. It is also possible that out of basin adult strays positively biased the adult estimate in 2012. Although the proportion of smolts that survive outside the Scott River watershed is largely driven by uncontrollable factors, it is important to track this survival metric as it provides a deeper understanding of adult return trend information.

Table 5. Coho Salmon smolt outmigrant abundance point estimates, age 2 and age 3 Coho Salmon abundance estimates, and proportion of outmigrant smolts that returned by brood year for the Scott River, Brood Years 2004-2019.

Brood Year	Smolt Year	Smolt point Estimate	Age 3 Return Year	Age 2 Return	Age 3 Return	Age 2 and 3 Return	Percent smolt survival
2004	2006	95815	2007	0	1539	1539	1.61
2005	2007	3931	2008	83	60	143	3.64
2006	2008	1142	2009	3	77	80	7.01
2007	2009	73232	2010	4	913	917	1.25
2008	2010	3257	2011	14	344	358	10.99
2009	2011	353	2012	11	188	197	56.37
2010	2012	63135	2013	13	2631	2644	4.19
2011	2013	9283	2014	121	383	504	5.43
2012	2014	6734	2015	102	188	290	4.31
2013	2015	8758	2016	24	226	250	2.85
2014	2016	3372	2017	0	368	364	10.91
2015	2017	N/A	2018	14	712	726	N/A
2016	2018	14628	2019	27	338	365	2.50
2017	2019	15707	2020	8	1664	1672	10.64
2018	2020	14628	2021	102	845	947	6.47
2019	2021	1762	2022	7	231	238	13.51

Analyzing comparisons of Coho Salmon smolt production estimates to estimated female adult Coho Salmon returns produces freshwater survival estimates in the form of Coho Salmon smolts produced per adult female return. For Brood Years 2007 through 2020, the number of Coho Salmon smolts produced per returning adult female has ranged from a low of 5.78 to a high of 101.78 and has averaged 57.46 (Table 6). Utilizing the average out of basin survival for yearling Coho Salmon of 6.1%, it requires a minimum of 32.8 yearlings to be produced per returning adult female to have zero population growth. Due to the difficulty in estimating abundance of adults and out-migrants at low abundance levels it is unclear if the smolts produced per adult female ratio generated for Brood Year 2009 is a result of decreased freshwater productivity or a result of sampling difficulty. As additional years of data become available the freshwater production of Coho Salmon in the Scott River can be further evaluated. The number of smolts produced per returning adult female by brood year is a direct measure of freshwater survival. For levels below carrying capacity, it can be stated that as the

number of smolts produced per returning adult female increases it can be inferred that in-river conditions for Coho Salmon are improving. Conversely as the number of smolts produced per returning adult female decreases, it can be inferred that in-river conditions for Coho Salmon are getting worse.

Table 6. Adult Coho Salmon estimate, Coho Salmon smolt production point estimate and ratio of Coho Salmon smolts produced per adult female return for the Scott River, Brood Years 2007-2015, 2016-2020.

Adult Year Brood Year	Adult Estimate	Adult Female Estimate	Smolt Year	Smolt point Estimate	Smolts produced per Female
2007	1622	860	2009	73,232	85.15
2008	63	32	2010	3,257	101.78
2009	81	41	2011	353	8.61
2010	927	640	2012	63,135	98.65
2011	355	170	2013	9,283	54.61
2012	201	86	2014	6,734	78.30
2013	2752	1514	2015	8,758	5.78
2014	485	179	2016	3,372	18.84
2015	212	170	2017	N/D	N/D
2016	226	175	2018	14,628	83.59
2017	382	174	2019	15,707	90.27
2018	739	422	2020	14,628	34.66
2019	346	195	2021	1,762	9.04
2020	1766	883	2022	68,616	77.71
				Average	57.46

When the number of returning adults is plotted against the number of Y+ Coho Salmon produced by brood year a positive relationship is generated (Figure 21). This positive relationship indicates that as adult spawners increase so does the resulting number of offspring. It is theorized that the extremely low production of offspring from the 2013 and 2019 brood years was the result of extreme drought conditions in the Scott River.




Figure 21. Number of returning adults and corresponding number of Y+ Coho Salmon produced by brood year in the Scott River, for Brood Years 2007-2014 and 2016-2020.

STEELHEAD

The number of returning adult steelhead has been monitored at the SRFCF since 2007. During the 2007 through 2009 seasons an unknown number of sub-adult steelhead may have been counted as adults. Starting in 2010, lines on the back of the video flume were set 16 inches (40.64 cm) apart to delineate sub-adults versus adults. From 2007 to 2022 the number of observed adult steelhead has ranged from a high of 917 to a low of 8 with an average of 235 (Figure 22). The run size of adult steelhead prior to 2007 is unknown, and with the addition of the counting facility, the ability to monitor this run has greatly improved. Although recent adult run size data is sparse on the Scott River, monitoring of the juvenile emigration has taken place since 2003. A large fraction of the adult steelhead migration occurs outside the operational window of the SRFCF. Therefore, the number of observed steelhead should be considered minimum number of returns and not basin estimates.




Figure 22. Number of observed steelhead >16” at the Scott River Fish Counting Facility from 2007 to 2022.

ACKNOWLEDGEMENTS

The California Department of Fish and Wildlife would like to thank Sport Fish Restoration Fund and the Pacific Coast Salmon Recovery Fund for their financial assistance in the operation of the SRFCF during 2022. The cooperative spawning ground surveys would not be possible without the assistance of the U.S. Fish and Wildlife Service Yreka office and the USFS. Additionally, this work would not be possible without the cooperation and participation of the Siskiyou Resource Conservation District, Karuk Tribe, Quartz Valley Tribe, United States Forest Service, Northern California Resource Center, California Conservation Corps in partnership with AmeriCorps, and Etna High School who assisted in completing spawning ground surveys on the Scott River. We would also like to express our appreciation to the various landowners who have provided permission to access the Scott River through their property. Additionally, we thank the SRCD for making their data available for this report.

References

- Bergman, J. M., R. M. Nielson, and A. Low. 2012. Central Valley in-river Chinook salmon escapement monitoring plan. Fisheries Branch Administrative Report Number: 2012-1. California Department of Fish and Game. Sacramento, CA.
- California Department of Fish and Wildlife (CDFW). 2021. Influence of Scott River in-stream flow on the distribution and migration timing of fall Chinook Salmon and Coho Salmon. California Department of Fish and Wildlife, Northern Region
- Giudice, D. and M. Knechtle. 2022. Shasta River Chinook and Coho observations in 2021-2022, Siskiyou County, CA. California Department of Fish and Wildlife. Available from 1625 South Main Street, Yreka, CA 96097
- Klamath River Technical Team (KRTAT). 2023. Klamath River Fall Chinook Salmon Age-Specific Escapement, River Harvest, and Run Size Estimates, 2022 Run.
- Voight, C. 2023. Personal communication. Siskiyou Resource Conservation District Etna, CA.
- Morrow, H. and A. Bachteler. 2022. 2022 Scott River Juvenile Salmonid Outmigrant Study. P1610529. California Department of Fish and Wildlife Available from California Department of Fish and Wildlife, 1625 South Main Street, Yreka, CA 96097
- Stevenson, David R., and Steven E. Campana. 1992. Otolith Microstructure Examination and Analysis. Department of Marine Resources, Marine Resources Laboratory, McKown Point, West Boothbay Harbor, ME 04575.
- Williams T. et al. 2008 Framework for assessing viability of threatened Coho salmon in the southern Oregon/northern California coast evolutionarily significant unit. NOAA-TM-NMFS-SWFSC-432.

